
UTAH - EXTENDED WEEKLY INFLUENZA SUMMARY

MMWR Week 1 (1/1/06 - 1/7/06) - Posted 1/11/06

WEEKLY MAP* OF UTAH INFLUENZA ACTIVITY



^{*} No Report/ No Activity * * Sentinel sites with regular ILI data collection are only located in the following counties: Cache, Davis, Salt Lake, Sanpete, Sevier, Tooele, Utah, Washington, and Weber.

Influenza-Associated Hospitalizations by Health District and County, and by Influenza Type - Utah 2005-2006 Week Ending: 1/07/2006 **Cumulative Cases** (MMWR Week 1) 09/03/2005 - Present Percent Total Type A Type B **Percent Total** Type A Type B **Health District** County **Percent Percent** *Unknown **Cases Percent Percent** *Unknown Cases 0 **Bear River** Box Elder 0.0 0.0 0.0 0 0.0 0.0 0 0.0 0.0 100.0 3 24 0.0 76.5 17 Cache Rich 0.0 0.0 0.0 0 0 0.0 0.0 0 Central 0.0 0 100 2 0.0 0.0 0.0 0.0 Juab 0 100 Millard 0.0 0.0 0.0 0.0 0.0 1 0 0.0 Piute 0.0 0.0 0.0 0 0.0 0 0.0 0.0 0.0 0 100 0.0 0.0 3 Sanpete 100.0 0.0 2 100 0.0 0.0 4 Sevier 0.0 Wayne 0.0 0.0 0.0 0 100 0.0 0.0 Davis 66.7 16.7 16.7 6 88 3.8 3.8 26 Davis Salt Lake 100.0 100.0 23 90 0.0 9.7 93 Salt Lake 0.0 100.0 1 Southeast Carbon 0.0 0.0 0.0 0.0 0.0 0.0 **Emery** 0.0 0.0 0.0 0.0 100.0 0.0 0.0 1 0.0 0.0 0 Grand 0.0 0.0 0.0 0.0 0.0 100.0 0.0 0.0 0.0 0.0 0.0 0.0 3 San Juan 0.0 0 Southwest Beaver 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0 Garfield 0.0 0.0 0.0 0.0 0.0 0.0 Iron 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0 0.0 0.0 0.0 0.0 100.0 0.0 0.0 1 Kane 27 0.0 Washington 100.0 0.0 0.0 4.0 100.0 0.0 Summit Summit 0.0 0.0 0.0 0.0 0.0 0 0.0 0.0 Tooele Tooele 0.0 0.0 0.0 0.0 100.0 0.0 0.0 4 **TriCounty** 0.0 0.0 0.0 100.0 0.0 0.0 1 0.0 Daggett Duchesne 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0 Uintah 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0 Utah 100.0 13.0 92.3 7.7 39 Utah 0.0 0.0 0.0 Wasatch Wasatch 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0 WeberMorgan Morgan 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0 75.0 25.0 0.0 4.0 93.5 6.5 0.0 31 Weber 86.8 State Total 8.9 56.0 2.3 10.5 257 87.5 3.6


NOTE: Reported influenza cases are collected to detect onset of influenza epidemics,


but account for only a small proportion of all persons who become ill from influenza.

Total Influenza-Associated Hospitalizations by Age Group, 2005-2006								
Age Group	Number of	Percentage of	Rate Per 100,000					
	Hospitalizations	Cases by Age	Person-Years					
< 1 year	19	11.2%	36.97					
1-4 years	13	7.7%	6.55					
5-14 years	7	4.1%	1.67					
15-24 years	2	1.2%	0.48					
25-34 years	8	4.7%	1.81					
35-44 years	12	7.1%	3.83					
45-54 years	9	5.3%	3.13					
55-64 years	15	8.9%	7.93					
65-74 years	22	13.0%	19.60					
75-84 years	33	19.5%	44.84					
85+ years	29	17.2%	108.49					
All ages*	169	100.0%	10.16					

^{*} Age data missing for 87 cases.

^{*} Unknown tests do not differentiate between influenza A and influenza B








Weekly Percentage of Visits to Utah Sentinel Providers that were Due to Influenza-Like Illness (ILI) by Age, Health District, and County - Utah 2005 - 2006


Health District	County	Age-Specific ILI Percentage			All Ages ILI	# of Sites	
		0 - 4	5 - 24	25 -64	65+	Percentage	Reporting
Bear River	Box Elder	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Cache	0.0%	0.0%	0.0%	0.0%	9.9%	1
	Rich	0.0%	0.0%	0.0%	0.0%	0.0%	0
Central	Juab	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Millard	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Piute	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Sanpete	49.8%	7.5%	11.9%	2.5%	13.4%	1
	Sevier	0.0%	0.0%	0.0%	0.0%	2.4%	1
	Wayne	0.0%	0.0%	0.0%	0.0%	0.0%	0
Davis	Davis	3.2%	2.3%	5.8%	0.0%	3.5%	2
Salt Lake	Salt Lake	5.7%	6.1%	3.9%	0.9%	3.9%	6
Southeast	Carbon	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Emery	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Grand	0.0%	0.0%	0.0%	0.0%	0.0%	0
	San Juan	0.0%	0.0%	0.0%	0.0%	0.0%	0
Southwest	Beaver	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Garfield	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Iron	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Kane	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Washington	0.0%	0.0%	0.0%	0.0%	0.0%	0
Summit	Summit	0.0%	0.0%	0.0%	0.0%	0.0%	0
Tooele	Tooele	6.7%	4.1%	0.5%	0.0%	1.5%	1
TriCounty	Daggett	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Duchesne	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Uintah	0.0%	0.0%	0.0%	0.0%	0.0%	0
Utah	Utah	5.8%	6.5%	10.6%	30.4%	6.1%	3
Wasatch	Wasatch	0.0%	0.0%	0.0%	0.0%	0.0%	0
WeberMorgan	Morgan	0.0%	0.0%	0.0%	0.0%	0.0%	0
	Weber	0.0%	0.0%	0.0%	0.0%	1.8%	1
Total		6.5%	5.7%	5.0%	2.6%	4.6%	16

Note: Some counties have not yet or do not report age-specific ILI percentages information. Age-specific data will be posted when available.









School Absenteeism Rates by County and Type of Absence Week Ending 1/7/06 (MMWR Week 1)								
Health District	County	All Absences Rate (absences/100 student-days)	Sick Absences Rate (absences/100 student-days)	ILI Absences Rate (absences/100 student-days)	# of Schools Reporting			
Bear River	Box Elder	0.0	0.0	0.0	0			
	Cache	0.0	0.0	0.0	0			
	Rich	0.0	0.0	0.0	0			
Central	Juab	0.0	0.0	0.0	0			
	Millard	0.0	0.0	0.0	0			
	Piute	0.0	0.0	0.0	0			
	Sanpete	7.0	2.6	0.4	3			
	Sevier	0.0	0.0	0.0	0			
	Wayne	0.0	0.0	0.0	0			
Davis	Davis	5.3	2.1	0.9	4			
Salt Lake	Salt Lake	6.1	0.0	0.0	6			
Southeast	Carbon	0.0	0.0	0.0	0			
	Emery	0.0	0.0	0.0	0			
	Grand	0.0	0.0	0.0	0			
	San Juan	0.0	0.0	0.0	0			
Southwest	Beaver	0.0	0.0	0.0	0			
	Garfield	0.0	0.0	0.0	0			
	Iron	0.0	0.0	0.0	0			
	Kane	0.0	0.0	0.0	0			
	Washington	0.0	0.0	0.0	0			
Summit	Summit	0.0	0.0	0.0	0			
Tooele	Tooele	1.9	0.0	0.0	12			
TriCounty	Daggett	0.0	0.0	0.0	0			
	Duchesne	0.0	0.0	0.0	0			
	Uintah	0.0	0.0	0.0	0			
Utah	Utah	1.9	0.0	1.3	2			
Wasatch	Wasatch	0.0	0.0	0.0	0			
WeberMorgan	Morgan	0.0	0.0	0.0	0			
	Weber	11.6	1.8	0.1	1			
Total		3.8	2.2	0.8	28			

UTAH - WEEKLY INFLUENZA SUMMARY

Influenza Activity in Utah:

While influenza activity appears to have peaked, influenza continues to have active circulation in Utah. The number of visits for influenza-like illness continues to be above state threshold levels. Student absences are also above threshold levels. Additionally, 257 hospitalizations associated with influenza have been reported to the UDOH this season. Most of these hospitalizations have occurred in individuals with high-risk factors and have been associated with Type A influenza.

As of 1/11/2006, no pediatric influenza-associated deaths have been reported in Utah.

Click on the map to access national influenza surveillance data. Please note, states determine for themselves how to estimate weekly influenza activity. The classification system used in Utah primarily serves to provide a spatial description of how influenza is circulating in the state.



INFLUENZA PREVENTION

How to protect yourself and others:

- 1. Utahns should get an influenza vaccine.
- **Contact your provider or the Immunization Hotline (1-800-275-0659) to locate an influenza shot.
- 2. All Utahns should practice "respiratory etiquette" to reduce the spread of respiratory diseases.
- ** Stay away from other people when you are sick.
- ** Cover your mouth and nose with a disposable tissue when you cough or sneeze. Throw the tissue away.
- ** Wash your hands.
- 3. We urge schools and employers to be supportive of these preventive measures.